

# NAMP Young Investigators Series: Environmental Radiochemistry

Dustin May  
Brennen Ferguson  
Frances Zengotita



National Analytical Management Program (NAMP)  
U.S. Department of Energy Carlsbad Field Office

TRAINING AND EDUCATION SUBCOMMITTEE



# Meet the Presenters...

**Dustin May** is a Ph.D. candidate in Human Toxicology at the University of Iowa working with Michael Schultz. He is an NRC Radiochemistry Fellow studying naturally occurring radioactivity in Iowa groundwater and its impact to public health, specifically focused on less well-understood radionuclides, polonium-210 and lead-210. He is also the manager of the State Hygienic Laboratory's radiochemistry department. May earned a B.S. in Chemistry also from the University of Iowa in 2006. He expects to graduate in December 2018.

Contact Information: [dustin-may@uiowa.edu](mailto:dustin-may@uiowa.edu)

Presentation: **Polonium-210 and lead-210 in groundwater**



# Meet the Presenters...

**Brennan Ferguson** is a Ph.D. student in Environmental Engineering and Earth Sciences at Clemson University working with Brian Powell. She is doing multi-scale experimental research on organic ligand facilitated dissolution of uranyl-phosphate minerals as part of a DOE Established Program to Stimulate Competitive Research (EPSCoR) task force. Ferguson has a B.A. in Chemistry and Environmental Studies from Alfred University in 2015 and expects to complete her Ph.D. work in December 2020.

Contact Information: [bofergu@g.clemson.edu](mailto:bofergu@g.clemson.edu)

Presentation: **Uranyl-phosphate mineral dissolution in the presence of organic ligands**



## Meet the Presenters...

**Frances Zengotita** is an undergraduate student pursuing a dual degree in Chemistry (B.S.) and English (B.A.) at Florida International University. She is a Department of Energy Fellow, a former Nuclear Regulatory Commission Scholar and a Ronald E. McNair Fellow under the mentorship of Dr. Hilary Emerson. Her research focuses on laboratory experiments to update risk assessment models on the behavior of the actinide series elements under high ionic strength conditions relevant to the WIPP. Zengotita expects to graduate in Spring 2019.



Contact Information: [fzeng002@fiu.edu](mailto:fzeng002@fiu.edu)

Presentation: **Nd<sup>3+</sup> Sorption to WIPP-Relevant Minerals**

**FIU**

Applied Research  
Center

# Polonium-210 and lead-210 in groundwater

Dustin May  
University of Iowa




National Analytical Management Program (NAMP)  
U.S. Department of Energy Carlsbad Field Office

TRAINING AND EDUCATION SUBCOMMITTEE




# Terrestrial NORM is Present in Groundwater


- Naturally Occurring Radioactive Material (NORM)
- Two primordial decay series of concern
  - Uranium series
  - Thorium series
- Persistent in environment
  - Parent radionuclide half-lives are billions of years



# Groundwater is the major source of drinking water in Iowa



Drinking Water Supplies by Source



Source: Iowa Geological Survey (Unpublished data)

# Some NORM is Regulated in Public Drinking Water

- Safe Drinking Water Act (SDWA) regulates contaminants in public drinking water
- Maximum Contaminant Level (MCL) for NORM based on excessive lifetime cancer risk
- $^{210}\text{Po}$  &  $^{210}\text{Pb}$  are not regulated directly
- Private drinking water wells are not regulated

Nuclide	MCL	Excessive Lifetime Cancer Risk
Radium-226 <sup>a</sup>	5 pCi/L (185 mBq/L)	$6.78 \times 10^{-5}$
Radium-228 <sup>a</sup>	5 pCi/L (185 mBq/L)	$1.89 \times 10^{-4}$
Uranium <sup>b</sup>	30 $\mu\text{g/L}$ (740 mBq/L)	$4.49 \times 10^{-5}$
Gross Alpha Particles	15 pCi/L (555 mBq/L)	N/A
Beta/Photon Emitters	4 mrem/year	N/A

<sup>a</sup> Enforced as Radium-226 + Radium-228

<sup>b</sup> Risk calculated based on assumed natural activity ratio (U-234:U238:U235, 1:1:0.046)

# Polonium-210 & Lead-210 Are Hazardous

## Po-210

- Alpha-emitter
- 138 day half-life
- Can exist as inorganic ions or organometallic complexes
  - Organometallic complexes more retained, ~50%
  - Po-210 is often found in a colloidal form in water
- Accumulates in reticuloendothelial tissues and erythrocytes
- Risk-based MCL = 1.1 pCi/L

## Pb-210

- Beta-emitting radionuclide
- 22.2 year half-life
  - Decays to Bi-210 and then Po-210
- Retained by body from 5-60%
  - Dependent on age and diet
  - Younger, higher retention
  - Fasted, higher retention
- Accumulates in erythrocytes and bone
- Risk-based MCL = 2.2 pCi/L

# Limited studies performed on $^{210}\text{Po}$

- Some studies have been done on  $^{210}\text{Po}$  in groundwater

- Seiler, et al. - Nevada
- Harada, et al. – Florida
- Ruberu, et al. – California
- Jacobus, et al. – Minnesota

- Studies found  $^{210}\text{Po}$  in groundwater
  - Limited by scope, geographic focus
  - Areas very different geologically



# Polonium-210 and lead-210 pose unknown risks



- Lack of wide-scale data examining  $^{210}\text{Po}$  and  $^{210}\text{Pb}$  in groundwater
- Alpha and beta MCLs too high to address potential risk
- Prevalence of NORM in private wells is unknown
- Higher potential carcinogenicity of  $^{210}\text{Po}$  and  $^{210}\text{Pb}$  relative to other NORM

# Aim of these studies

- Identify drinking water resource types (private wells; surface supply; deep aquifer) and hydrogeological character (well depth/age, geological deposit, aquifer) in which elevated levels of regulated NORM and unregulated  $^{210}\text{Po}$  are found.




# Study Design




- Samples collected from all over the state of Iowa
  - 119 samples from public water supplies
  - 39 samples private wells
- Samples collected in representative manner
  - Based on drinking water withdrawals
- Analyzed for uranium-series radionuclides, various water quality parameters
  - TSS, TDS, Chloride, Alkalinity, Nitrate, Ammonia, Phosphorus

# FY16 Public Well Sites



# FY18 Public Wells Sites




# Methods and Quality Control

- Analyses performed using a variety of methods
  - EPA 900.0 (Gross Alpha/Beta)
  - EPA 903.0 ( $^{226}\text{Ra}$ )
  - Eichrom methods (U,  $^{210}\text{Pb}$ )
  - In-house adapted method for  $^{210}\text{Po}$  analysis
    - $\text{MnO}_2$  pre-concentration, auto-deposition on nickel disc
- Method performance was verified using appropriate quality controls
  - Laboratory control samples, matrix spikes, duplicates, reagent blanks




# Radiological Parameters Vary Across Aquifers







# Po-210 Distribution is Different from Gross Alpha




# Elevated $^{210}\text{Po}$ Well Below Alpha MCL




# Gross Beta Relationship with Well Depth is Less Clear



# $^{210}\text{Pb}$ Not Related to Well Depth



# Highest $^{210}\text{Pb}$ Found at Low Beta Levels



# Conclusions

- The Dakota aquifer appears different
  - Higher  $^{210}\text{Po}$ , U activities
  - Mean gross alpha activity higher than beta activity
- Gross alpha/beta screening is inadequate for  $^{210}\text{Pb}$ ,  $^{210}\text{Po}$ 
  - $^{210}\text{Po}$  shows significant negative correlation with well depth, higher at lower gross alpha activities
  - $^{210}\text{Pb}$  also higher at lower gross beta activities
- All  $^{210}\text{Po}$ ,  $^{210}\text{Pb}$  less than the theoretical MCLs
  - But,  $\sim 7\%$  of  $^{210}\text{Po} > \frac{1}{2}$  MCL

# Future Work

- **Private Well Study**
  - Analyses to be completed this summer
  - Comparison to active public water supplies
- **Follow up on Dakota aquifer anomalies**
  - How is this aquifer different?
  - Are these differences widespread? reproducible?

# Acknowledgements

- Thank you to:
  - Michael Schultz, PhD
  - Claire Hruby, PhD
  - Michael Wichman, PhD
  - Amanda Carl
  - Lindsey Tirado
- Iowa DNR
- State Hygienic Laboratory
- CHEEC
- NRC



University of Iowa



Radiochemistry



THE UNIVERSITY  
OF IOWA

Center for Health Effects  
of Environmental Contamination



Interdisciplinary Graduate Program in

# HUMANTOXICOLOGY

# References

- Cutler Prior, J.B., Janice; Howes, Mary; Libra, Robert; VanDorpe, Paul, *Iowa's Groundwater Basics*. 2003, Iowa Department of Natural Resources: Iowa City, IA.
- U.S. EPA. *Information about Public Water Systems*. March 8th, 2017 November 19th, 2017]; Available from: <https://www.epa.gov/dwreginfo/information-about-public-water-systems>.
- Eckerman, K.F., et al., *Federal Guidance Report No. 13: Cancer Risk Coefficients for Environmental Exposure to Radionuclides*. 1999: Washington D.C.
- Thomas, P.A., et al., *Human absorption and retention of polonium-210 from caribou meat*. *Radiat Prot Dosimetry*, 2001. **97**(3): p. 241-50.
- Agency for Toxic Substances and Disease Registry, *Toxicological Profile for Lead*. 2007, U.S. Public Health Service: Washington, D.C.
- Rabinowitz, M.B., G.W. Wetherill, and J.D. Kopple, *Kinetic analysis of lead metabolism in healthy humans*. *Journal of Clinical Investigation*, 1976. **58**(2): p. 260-270.
- Stannard, J.N., *Distribution and Excretion of Polonium-210. III. Long-Term Retention and Distribution in the Rat*. *Radiation Research Supplement*, 1964. **5**: p. 67-79.
- Ruberu, S.R., Y.G. Liu, and S.K. Perera, *Occurrence and distribution of 210Pb and 210Po in selected California groundwater wells*. *Health Phys*, 2007. **92**(5): p. 432-41.
- Jacobus, J., et al., *Polonium-210 Occurrence in Minnesota's Aquifers: A Pilot Study*. 2014, Minnesota Department of Public Health: St Paul, MN.
- Seiler, R.L., et al., *Biogeochemical factors affecting the presence of 210Po in groundwater*. *Applied Geochemistry*, 2011. **26**(4): p. 526-539.
- Harada, K., et al., *Polonium in Florida groundwater and its possible relationship to the sulfur cycle and bacteria*. *Geochimica et Cosmochimica Acta*, 1989. **53**(1): p. 143-150.
- Eckerman, K.F., et al., *Federal Guidance Report No. 13: Cancer Risk Coefficients for Environmental Exposure to Radionuclides*. 1999: Washington D.C.

# Uranyl phosphate mineral dissolution in the presence of organic ligands

Brennan Ferguson  
Clemson University



National Analytical Management Program (NAMP)  
U.S. Department of Energy Carlsbad Field Office

TRAINING AND EDUCATION SUBCOMMITTEE





# Dangers of Uranium

- Uranium contaminates 120 sites in 36 states
  - Mining and milling
  - Nuclear Energy and Weapons
  - Natural Processes
- Public health concerns
- $\text{UO}_2^{+2}$  - soluble




# System of Interest

- **Uranyl-phosphate:**
  - Stable under oxidizing conditions
  - Precipitates faster and less soluble than non-phosphate minerals
  - Can be dissolved by organic ligands: citrate, EDTA, acetic acid, bicarbonate
- **Citrate:**
  - Produced by plant roots to mobilize nutrients
  - Increases uranium accumulation in plants



# Chernikovite: A Uranyl-Phosphate Precipitate


- Chemical Formula:  
 $(\text{H}_3\text{O})(\text{UO}_2)(\text{PO}_4)\cdot 3\text{H}_2\text{O}$



Mineral	Chemical Formula	Log $K_{sp}$
Chernikovite	$(\text{H}_3\text{O})(\text{UO}_2)(\text{HPO}_4)\cdot 3\text{H}_2\text{O}$	-22.73
Autunite	$\text{Ca}[(\text{UO}_2)(\text{PO}_4)]_2\cdot 4\text{H}_2\text{O}$	-48.36
Uranyl Orthophosphate	$(\text{UO}_2)_3(\text{PO}_4)_2\cdot 4\text{H}_2\text{O}$	-49.36

# Uranyl-Phosphate Precipitate

- **Synthesis:**
  - Mix  $\text{UO}_2(\text{NO}_3)_2$  and  $\text{Na}_2\text{HPO}_4$
  - Yellow precipitate forms immediately
  - For one experiment, added citrate to  $\text{Na}_2\text{HPO}_4$  solution
- **Characterization:**
  - X-ray diffraction – chernikovite
  - Scanning electron microscope (SEM) with electron dispersive spectroscopy (EDS)




	Wt%	$\sigma$
U	52.7	0.2
O	24.7	0.2
C	15.8	0.2
P	5.5	0.1
Na	1.4	0.1

# Motivation

- **Specific:** to examine the kinetics and extent of citrate facilitated dissolution of chernikovite


# Motivation

- Specific: to examine the kinetics and extent of citrate facilitated dissolution of chernikovite ( $\text{UO}_2\text{HPO}_4 \cdot 4\text{H}_2\text{O}$ )
- **Expanding work and collaboration:** micromodels, plant uptake, and field scale studies



# Motivation

- **Specific:** to examine the kinetics and extent of citrate facilitated dissolution of chernikovite ( $\text{UO}_2\text{HPO}_4 \cdot 4\text{H}_2\text{O}$ )
- **Expanding work and collaboration:** micromodels, plant uptake and field scale studies
- **Broader Impact:** to gain a better understanding of the fate of uranyl-phosphate precipitates in the environment




# Research Questions

1. What is the relationship between **citrate** concentrations and **dissolved uranium** from chernikovite?
2. When does the reaction reach **steady state**?
3. Do **citrate and phosphorous** in solution **sorb** to the chernikovite surface?
4. Does the citrate affect the **remaining solid**?

# Experiments


- Batch dissolution – varying citrate concentrations
- Continuous stirred-tank reactor (CSTR) – varying citrate concentrations and flow rate
- Isotope sorption study with  $^{32}\text{P}$  and  $^{14}\text{C}$
- Post dissolution analysis of precipitate – Raman, theoretical simulations, x-ray diffraction (XRD)

# Dissolved Uranium Increases with Citrate




**Dissolution of chernikovite in a batch experiments for 12 hours with 12 different concentrations of citrate at pH 4**

# Continuous Stirred-Tank Reactor (CSTR) Experiments




- Enable measurements of rapid reactions
- Minimizes transport phenomena
- Experiments
  - Varied citrate concentration: no citrate, 0.1 mM, 1 mM, and 10 mM
  - Pretreatment: Wash for 10 reactor volumes (RV) with 10 mM NaCl, citrate for 15 RV
  - Varied flow: Wash, 1 mL/min, 0.5 mL/min, 0.25 mL/min
  - Constant pH of ~4

# Reaching Steady State - CSTR Results




Dissolution of chernikovite in a continuously stirred tank reactor with four different concentrations of citrate (denoted by marker color) and three different flow rates (denoted by marker shape) at pH 4

# Fast Citrate and Phosphorous Exchange




After 24 hours, a 0.1 mM citrate and chernikovite batch system was spiked with  $^{32}\text{P}$  and  $^{14}\text{C}$  labeled citrate.




# Raman - Structure Changes


Post CSTR Chernikovite and  
Theoretical Simulations



- $\text{PO}_4^{-3}$  stretching: 986 – 1020  $\text{cm}^{-1}$
- $\text{UO}_2^{+2}$  stretching: 821 – 867  $\text{cm}^{-1}$
- Downshift in  $\text{UO}_2^{+2}$  peak with increasing citrate concentration
- Theoretical simulation: 1:2 uranyl to citrate ratio with middle carboxylic group



# XRD - Structure Changes



# Research Questions → Research Findings

1. What is the relationship between citrate concentrations and dissolved uranium from chernikovite?
  - a) The presence of citrate increases the concentration of mobile uranium.
  - b) Citrate surface saturation or reprecipitation hinders the dissolution.
2. When does the reaction reach steady state?

Steady state dissolution of chernikovite was achieved within 30 min but only after the precipitate was pre-treated with citrate to achieve a more uniform chemical environment for dissolution.

# Research Questions → Research Findings

3. Do citrate and phosphorous in solution sorb to the chernikovite surface?

Phosphorous and citrate in solution exchanges quickly with the phosphorous and citrate on the mineral surface, with possible re-precipitation of phosphate.

4. Does the citrate affect the remaining solid?

Citrate alters the structure of the remaining chernikovite.

# Broader Impact

- Uranyl-phosphate minerals may not be as immobile as originally thought
  - Soil rich in organic matter
  - Bacteria processes
  - Plant exudates

# References

- National Research Council 2010 Science and Technology for DOE site Cleanup: Workshop summary
- National Research Council (NRC) (2008), Review of the Toxicologic and Radiologic Risks to Military Personnel From Exposures to Depleted Uranium During and After Combat, Natl. Acad. Press, Washington, D. C.
- Palmisano, A., Hazen, T., 2003. Bioremediation of Metals and Radionuclides: What It is and How It Works, second ed. LBNLe42595-(2003)
- Image slide 2: [https://en.wikipedia.org/wiki/Ionizing\\_radiation](https://en.wikipedia.org/wiki/Ionizing_radiation) (accessed April 18, 2016)
- Fanizza, M.F., et al. *Water Resources Research*, 2013. 49(2): p. 874-890.
- Sowder, A.G., et al. *Journal of Radioanalytical and Nuclear Chemistry*, 2001. 248: p. 517-524. Berto, S., et al. *Radiochim. Acta*, 2012. 100: p. 13-28.
- Gorman-Lewis, et al., *Environ. Sci. Technol.* 43 (2009) 7416-7422.
- Image slide 7: Powell et al. “Radionuclide Waste Disposal: Development of Multi-scale Experimental Modeling Capabilities.” EPSCor All Hands Meeting: Clemson, SC. July, 2015.
- Image slide 8: <http://ens-newswire.com/2013/02/23/six-storage-tanks-leaking-at-hanford-nuclear-site/> (accessed April 2016)

# Acknowledgements

- Dr. Brian Powell and the Powell research group
- Dr. Lindsay Shuller-Nickles
- Dr. Lawrence Murdoch
- Fengjiao Liu, Dr. Apparao Rao, Dr. Ramakrishna Podila for the Raman help
- Dr. Michael Trumm for the theoretical calculations

This material is based upon work supported by the U.S. Department of Energy Office of Science, Office of Basic Energy Sciences and Office of Biological and Environmental Research under Award Number DE-SC-00012530



U.S. DEPARTMENT OF  
**ENERGY**

Office of  
Science

# $\text{Nd}^{3+}$ SORPTION TO WIPP-RELEVANT MINERALS

FRANCES ZENGOTITA

FLORIDA INTERNATIONAL UNIVERSITY




National Analytical Management Program (NAMP)  
U.S. Department of Energy Carlsbad Field Office

TRAINING AND EDUCATION SUBCOMMITTEE



# Nuclear Waste clean-up

- Large amounts of radioactive waste began to accumulate in the 1940s as a byproduct from WWII and then the Cold War.
- A deep geologic repository was ultimately recommended as the best option for long-term disposal by the National Academy of Sciences.
- During the 1960s, the Chihuahuan desert of southeastern New Mexico was chosen to be an appropriate site.



# Repository Storage for Transuranic Waste




Figure 2. TRU Defense Waste Generating and Storage Sites

## What is TRU Waste?

TRU Waste (Transuranic Waste) – radioactive waste containing  $>100$  nCi/g of alpha from elements greater than or equal to atomic # 92 (U)


## How much waste is at the WIPP\*?

- 90,600 m<sup>3</sup> CHTRU
- 360 m<sup>3</sup> RHTRU

\*received through 11,900 shipments as of 2014


# Waste Isolation Pilot Plant (WIPP)

- The WIPP is a deep geologic repository for the permanent disposal of radioactive waste that is a byproduct from the nation's nuclear defense program.
- Only plant in the world to handle and permanently dispose of transuranic waste.



# WIPP Release Scenario

- In the most likely WIPP Release scenario, human intrusion can lead to direct and/or long-term brine release (US DOE 1995, US DOE 1996, Perkins et al., 1999).
- Once the brine is released, it may proceed through the Rustler formation (the most transmissive layer) and pose a potential threat to the environment (Perkins et al., 1999)



## Lack of Experimental Sorption Data for Trivalent Actinides/Lanthanides for Dolomite

Previously,  $\text{Log}K_d$ 's for dolomite highly variable, reported from 3.4 – 6

- **Brady et al. 1999**: limited residence time reactor measured  $K_d$ 's, exceeded Nd solubility for pH 6 – 8, limited pH range for Am(III) (pH 3 – 6), limited conditions (0.05 and 0.5 M NaCl)
- **Perkins et al. 1999**: intact-core experiments, no breakthrough recorded after many months of injection, possible precipitation of Am(III)
- **Brush and Storz 1996**: batch sorption experiments in brines, no pH reported for Am(III)  $K_d$ 's, limited Nd(III)  $K_d$ 's for atmospheric  $\text{CO}_2$  in 0.05 M NaCl


# OBJECTIVES

- The focus of this research is to investigate sorption processes of neodymium to dolomite minerals from the Rustler formation above the WIPP repository.
  - **Relevance:** These data will decrease uncertainty in the WIPP risk assessment models by decreasing the current range of sorption parameters for the trivalent actinides.
  - Investigated the effects of
    1. **Ionic strength** (Borkowski *et al.*, 2009; Emerson *et al.*, 2018)
    2. **Ligands** (Emerson *et al.*, 2017-18; Roach *et al.*, 2008)
    3. **Microbes** (Zengotita *et al.*, 2017)

# Neodymium Background

## So why are we using Nd for this work?

- Nd<sup>3+</sup> exhibits similar chemical behavior to Am<sup>3+</sup> as it exists in the same oxidation state (Silva and Nitsche, 1995)
- Nd solubility is ~50 ppb at pH 8.5
  - Strong carbonate complexation occurs
  - Modeled in 0.1M solution
- Solubility in 5.0 M NaCl has been measured from 715 to 0.1 ppb ( $5 \times 10^{-6}$  to  $1 \times 10^{-9}$  M) at pH 8.5 in over and under-saturation experiments highlighting that we still do not fully understand the high IS system (Borkowski *et al.*, 2009).




Nd solubility in 0.1 M NaCl + atmospheric CO<sub>2</sub>

# Experimental Design: Mini Columns

## Initial Conditions

- 2.2 cm column with (1 cm of dolomite + fittings) (Top figure)
- Kohlen Pump (Bottom figure) pushes electrolyte background + Nd into mini-column
- Effluent is collected in 13x100 mm polystyrene tubes in Eldex fraction collector
- Experimental Design inspired by:
  - Dittrich, T.M., Ware, S.D., Reimus, P.W. (2016) Mini-columns for Conducting Breakthrough Experiments: Design and Construction, in: Technologies, U.S. DOE. Los Alamos National Laboratory, Los Alamos, NM. LA-UR-15-24392



# Experimental Design: Batch Experiments

- Initial Conditions
  - Variable concentrations of dolomite 0.5, 1.0, 5.0 g/L
  - Background electrolyte of 0.01 to 5 M total ionic strength [3 mM  $\text{NaHCO}_3$  + NaCl] + 20 pbb Nd + ligands
- Sampling Protocol
  - Variable time measurements: 15 minutes to 60 days
  - 15 minute settling, centrifugation, or filtration prior to sampling



End over end mixer with variable solid to liquid ratio batch sorption experiments

# High Ionic Strength Systems

## ➤ Why high Ionic Strength?

➤ The WIPP repository that contains the nuclear waste is a salty layer (up to 7.4 M).

➤ High ionic strength systems are not as well understood (Borkowski *et al.*, 2009).

## ➤ What is $pC_H$ ?

➤  $pC_H$  values normalize the ionic strength in the solution in terms of  $H^+$  ion concentration (Borkowski *et al.*, 2009).

➤ The pH measurement is not straightforward in high ionic strength solutions due to changes in *activity*.


## Relevant Equations

$$\text{Eqn. 1: } \Delta\text{pH} = 0.1868[I] + 0.073$$

$$\text{Eqn. 2: } \Delta\text{pH} = pC_H - \text{pH}$$

$$\text{Eqn. 3: } I = \frac{1}{2} \sum c_i z_i^2$$

# Gran Titrations at Variable IS




## LEGEND:

**Blue** Our data


**Navy** (Rai *et al.*, 1995)

## Results: Variable Ionic Strength




Sorption (as  $K_{d,con}$  in mL/g) with respect to ionic strength for 5 g/L dolomite and initially 20 ppb Nd (Emerson *et al.*, 2018)

## Results: Variable Ionic Strength



It is likely that the removal of Nd increases as ionic strength increases and affects mineral dissolution and precipitation processes. Previous research (Morse *et al.*, 2002; Lucero *et al.*, 1998) has shown that Nd is removed by (1) a fast adsorption process followed by (2) a slow co-precipitation process.

## Results: Variable Ionic Strength



**Why are the results from both experiments different?**

- Different removal processes in different systems due to the reaction times
- Two reactions are occurring in batch experiments (1) fast adsorption (2) slow process with co-precipitation
- Equilibrium is not reached in the mini-column due to the short reaction time, i.e. co-precipitation does not occur. (Emerson *et al.*, 2018)

# Biocolloid Transport of Nd

**Sub-objective:** To observe the behavior of *Chromohalobacter* and its effect on the mobility of Nd in the presence of dolomite minerals.

We considered:

- (1) Co-transport of Nd with microbes and
- (2) Re-mobilization of Nd adsorbed to dolomite by microbes (Zengotita *et al.*, 2017).

## **WHY CHROMOHALOBACTER?**

- Isolated from near the WIPP site,
- Can thrive in high salt concentrations which are relevant to WIPP-related conditions,
- Very little experimental data exists on the behavior of *Chromohalobacter* species for WIPP-related conditions





K.S.111  
Print Mag: 34100x @ 7.0 in  
14:15 07/01/10  
Microscopist: pcooke

500 nm  
HV: 80KV  
Direct Mag: 38000x  
X: 626.7 V, -449.4  
RMCU

(Dittrich, T.M., et al, 2017)

# Results: Co-transport of Nd with Biocolloids



## Results for Column with initial injection of *Chromohalobacter* + Nd + brine


- **Unfiltered (left):** Highlights good correlation of Nd and Microbes
- **Filtered (right):** Suggest majority of aqueous neodymium associated with microbes (<3% dissolved)
  - (Zengotita *et al.*, 2017)

### LEGEND:

- Blue Nd unfiltered recovery
- Red Nd filtered recovery
- Green microbe recovery

\*Note change in y-axis between results

# Results: Remobilization of Nd by Biocolloids



## Results for column with initial injection of Nd

- Strong sorption onto the dolomite
- Microbial recoveries show no association with Nd
- No remobilization

## Summary

- The microbes were unable to pull Nd off the dolomite.
  - (Zengotita *et al.*, 2017)

Results for column with initial injection of 20 ppb Nd + brine followed by injection of *Chromohalobacter* + brine only (only unfiltered data presented)


### LEGEND:

- Blue Nd unfiltered recovery
- Red Nd filtered recovery
- Green microbe recovery

# Effect of EDTA on Nd Transport

## Sub-objective:


The main focus of this task is to investigate the binding effect of the EDTA on Nd in the aqueous state and observe the potential removal of the contaminant on the surface of the dolomite.



## Why EDTA?

- EDTA concentrations in the WIPP could reach 0.3 mM!
- EDTA is used in nuclear decontamination chemistry because of its ability to bind radionuclides and remove them from contaminated surfaces (Roach *et al.*, 2008).
- It is persistent in the environment which makes it relevant in studies of radioactive waste disposal.
- Forms stable complexes with cationic metals (for example:  $\text{Nd}^{3+}$ )
  - (Rizkalla *et al.*, 1989)

# Results: Effect of EDTA on Nd Transport



## ➤ Initial Conditions

- 1000ppb Nd
- 5 g/L dolomite
- 5mg/L EDTA
- IS: 0.1M

\*30 & 60 day both at equilibrium

***So... what does this graph mean?***

- At 24 hours, very little sorption of Nd to dolomite with EDTA
- Around the 30 day period, the dolomite started to compete with the EDTA in respect to Nd sorption.
- The 60 day period demonstrates that the EDTA was able to complex the Nd and out-compete the dolomite

Note: The complexation that is adsorbing to the dolomite is the Nd-carbonate

# Conclusions

These experiments updated parameters for risk assessment models on the environmental behavior of the actinide series elements.

1. In the absence of ligands, strong sorption of Nd to dolomite occurs from 0.1 – 5.0 M IS.
2. Ionic strength has an impact on Nd removal processes in the absence of flow potentially due to its effect on mineral dissolution and co-precipitation processes.
3. Microbes can mobilize Nd if they interact with Nd first but cannot remobilize the Nd from dolomite. However, different microbes may interact more strongly.
4. EDTA reduces sorption to dolomite ( $K_{d,cond}$ ).

## ***Why is this relevant?***

- These data help us to better understand the potential for transport of the actinide elements from the WIPP and allow us to identify which parameters may have the greatest impact on mobility.
- The results suggest that the WIPP should consider the environmental conditions (based on the salt formation, high IS systems), presence of EDTA (found in storage and usage) and microbes. These data demonstrate that there are multiple potential transport vectors for actinides in the storage of nuclear waste.

# References

1. Morse, J.W., Arvidson, R.S. (2002) The dissolution kinetics of major sedimentary carbonate minerals. *Earth-Science Reviews*, **58**, 51-84
2. Morse, J.W., Mackenzie, F.T., 1990. *Geochemistry of Sedimentary Carbonates*. El Sevier, New York, NY.
3. Brady, P.V., Papenguth, H.W., Kelly, J.W. (1999) Metal sorption to dolomite surfaces. *Appl Geochem*, **14**, 569-579.
4. Keith-Roach, M.J. (2008). The speciation, stability, solubility and biodegradation of organic co-contaminant radionuclide complexes: a review. *The Science of the total environment*, *396* 1, 1-11.
5. Perkins, W.G., Lucero, D.A., Brown, G.O. (1999) Column Experiments for Radionuclide Adsorption Studies of the Culebra Dolomite: Retardation Parameter Estimation for Non-Eluted Actinide Species, in: Laboratories, S.N. (Ed.), Albuquerque, NM, p. 163.
6. Lucero, D.A., Heath, C., Brown, G. (1998) Laboratory column experiments for radionuclide adsorption studies of the Culebra Dolomite Member of the Rustler Formation. Sandia National Labs., Albuquerque, NM.
7. Dittrich, T.M., Ware, S.D., Reimus, P.W. (2016) Mini-columns for Conducting Breakthrough Experiments: Design and Construction, in: Technologies, U.S. DOE. Los Alamos National Laboratory, Los Alamos, NM. LA-UR-15-24392
8. Borkowski, Marian, Lucchini, Jean-Francois, Richmann, Michael K., & Reed, Donald T. *Actinide (III) solubility in WIPP Brine: data summary and recommendations*. United States. doi:10.2172/966984.
9. Zengotita, Frances, Emerson, Hilary Palmer, Dittrich, Timothy M., Swanson, Juliet S., & Reed, Donald T. *The Role of Chromohalobacter on Transport of Lanthanides and Cesium in the Dolomite Mineral System*. United States. doi:10.2172/1411368.
10. Emerson, H., Katsenovich, Y., Lagos., L (2018) Task 5 Remediation Research and Technical Support for WIPP, *Project Technical Plan*, Miami FL
11. Emerson, H., Zengotita, F., Richmann, M., Katsenovich, Y., Reed, D.T., Dittrich, T., (2018) Retention of neodymium by dolomite at variable ionic strength as probed by batch and column experiments. *Journal of Environmental Radioactivity*. Miami, Florida
12. Rai, D., Felmy, A.R., Juracich, S.P., & Rao, F. *Estimating the hydrogen ion concentration in concentrated NaCl and Na<sub>2</sub>SO<sub>4</sub> electrolytes*. United States. doi:10.2172/106419.

# Acknowledgements

**LANL Summer Mentors**

**Tim Dittrich, Donald Reed, and Juliet Swanson**

**FIU ARC Mentors**

**Hilary Emerson and  
Yelena Katsenovich**

**DOE-FIU Science and Technology Workforce Development Program**

**Sponsored by the U.S. Department of Energy, Office of Environmental  
Management, under Cooperative Agreement**

**#DE-EM0000598 (PI Dr. Lagos)**

Questions for our presenters??

A decorative graphic consisting of a solid green horizontal bar that spans the width of the slide. Below this bar, on the right side, there are several thin, parallel white lines that create a stepped or layered effect, extending further to the right.

# Upcoming Webinars

- Young Investigators Series: Separation Science
- Young Investigators Series: Radioecology
- Young Investigators Series: Novel Uranium Nanostructures

NAMP website <http://www.wipp.energy.gov/namp/>